BEST COPY AVAILABLE

139

R. D. O'Heill, Administrative Officer Division of Biology and Medicine March 9, 1953

Walter D. Claus, Chief Biophysics Branch, Division of Biology and Fedicine

MATERIAL FOR MONTELY PROCHESS REPORT - FEBRUARY 1953

R

Residual Environmental Contamination - Tembler/Snapper Data.

Observations on the intensity of radioactive fallout in areas near the Nevada Proving Ground have been continued since the test series in 1952 (Tumbler/Snapper). Calculations from the data observed at Groom and Lincoln mines, and the town of Pioche show that in January 1953, residual levels of activity should amount to about 0.5, 0.1 and 0.5 milliroentgens per hour, respectively. Actual observations, however, show that the radiation levels are approximately normal background, i.e., 0.02 - 0.03 milliroentgen per hour. It thus appears that in addition to radioactive decay, other factors such as weathering (scattering by winds, penetration into soil as a result of manow and rain), are effective in reducing residual environmental decage effects far below the levels which have been considered. It omental hazards due to radioactive fallout in populated areas near the nite have been overestimated.

Title of Project: "The Separation of the Proteins of Lymphoid Tissue and Their Theracterization as Regards Sadiation Sensitivity". (Unclassified) Professional Personnel: Engene L. Ress, Principal Investigator Institution: Sorthwestern University Medical School

Pr. Wess and associates, investigating methods for separation of various components of lymphoid tissue, have isolated a protein in the cytoplasmic nucleoprotein fraction which has unusual sensitivity of ultraviolet light and to alkalinity. Since lymphoid tissue is known to be highly sensitive to radiation damage this finding may prove to be significant. Additional work is needed to further purify and characterize this component.

Meteorological Program at Brookhaven, (Unclassified)

A meteorological research program has been approved recently at Brookhaven National Laboratory to study the phenomena of turbulent diffusion and air movements at both high and low levels. The Laboratory has unique facilities and excellent equipment which can be utilized for such studies. In addition, an experienced staff is available which is thoroughly familiar with meteorological phenomena of this type, and is also cognizant of the problems associated with Commission operations.

The effect of weather on Commission activities introduces a variety of problems of atmospheric pollution, radioactive affluents, fallout

OFFICE	BMBP		ADDALISTATION O	- RAAMAAFRAFMT	1 1
CHONAME	CLAUS me		UKGARIZATION &	k IMANAGEMENI /	41
	3-9-53	क्षण १५५ के दसन कर्ण स	CO TOTAL COMPANY	Ž	(
Form AEC	-318	U. S. GOVERNMENT PRINTING OFFICE	10 OBIGINA	*	

DOS ARCHIVES

phenomena, and dispersion of bomb debris. Under the field investigations which are planned, emphasis will be placed on atmospheric turbulence and the measurement of meteorological variables using radioactive isotopes. The development of such data has basic implications in the study of industrial hygiene, waste disposal methods, and environment as related to the production activities of the Commission.

mc (rewritten by F. Laner)

US DOE ARCHIVES
326 U.S. ATOMIC ENERGY
COMMISSION

RG_	DOE	HISTORIAN	(DBM)		
Colle	ction _	1/32			
Box _ 3363					
Folde	r <u>#</u> .	23			

OFFICE ▶						
SURNAME >						
DATE >			1,1 1,100,100			
Form AEC	·	U, S.	GOVERNMENT PRINTING OFFICE	16-62761-2	*	